

PCT/GB 2004 / 0 0 2 8 2

30 JUNE 2004

REC'D 15 JUL 2004

WIPO

PCT

PA 1104699

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

June 21, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/484,032

FILING DATE: June 30, 2003

PRIORITY

DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



L. Edele

L. EDELEN
Certifying Officer

PATENT APPLICATION SERIAL NO. _____

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE
FEE RECORD SHEET

07/03/2003 HTECKLU1 00000076 250166 60484032

01 FC:1005 160.00 DR

PTO-1556
(5/87)

U.S. Government Printing Office: 2001 — 481-697/58173

PROVISIONAL APPLICATION FOR PATENT COVER SHEET
 This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. ET971611288US

INVENTOR(S)			
Given Name (first and middle (if any)) Irene	Family Name or Surname Karantzeni	Residence (City and either State or Foreign Country) Waltham, Massachusetts	
<input checked="" type="checkbox"/> Additional Inventors are being named on the _____ separately numbered sheets attached hereto			
TITLE OF THE INVENTION (600 characters max) ISOTHERMAL TITRATION CALORIMETRY ASSAYS			
Direct all correspondence to:			
CORRESPONDENCE ADDRESS			
<input checked="" type="checkbox"/> Customer Number <div style="border: 1px solid black; width: 100px; height: 20px; margin: 5px; text-align: center;">22466</div>	<div style="border: 1px solid black; width: 200px; height: 40px; margin: 5px;"></div> Place Customer Number Bar Code Label here		
OR Type Customer Number here			
<input type="checkbox"/> Firm or Individual Name			
Address			
City			
Country		State	ZIP
		Telephone	Fax
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification Number of Pages	<input type="checkbox"/> CD(s), Number	<input type="checkbox"/> Other (specify)	
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets	<input checked="" type="checkbox"/> Other (specify)		
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.78			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT			
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.			
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.			
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:			
Payment by credit card. Form PTO-2038 is attached.			
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.			
<input checked="" type="checkbox"/> No.			
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are:			

Respectfully submitted,

(Page 1 of 2)

SIGNATURE *Robin S. Quartin*
 TYPED or PRINTED NAME Robin S. Quartin
 TELEPHONE 302-885-9129

Date 06/30/03

REGISTRATION NO.
 (If appropriate)
 Docket Number:

45,028

100997-1/US

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.61. This information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 6 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22315-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22315-1450.

If you need assistance in completing the form, call 1-800-PTO-9169 and select option 2.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Approved for use through 04/30/2003. CMB 0681-0002
U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

FEE TRANSMITTAL for FY 2003

Effective 01/01/2003, Patent fees are subject to annual revision.

☐ Applicant claims small entity status. See 37 CFR 1.27

TOTAL AMOUNT OF PAYMENT (\$)**160.00**

Complete If Known

Application Number	Not yet assigned
Filing Date	June 30, 2003
First Named Inventor	Karantzeni et al.
Examiner Name	Not yet assigned
Art Unit	Not yet assigned
Attorney Docket No.	100997-1/OS

METHOD OF PAYMENT (check all that apply)

☐ Check ☐ Credit card ☐ Money Order ☐ Other ☐ None

☒ Deposit Account:

Deposit Account Number: **26-0166**

Deposit Account Name: **AstraZeneca**

The Commissioner is authorized to (check all that apply):
☒ Charge fee(s) indicated below ☐ Credit any overpayments
☒ Charge any additional fee(s) during the pendency of this application
☐ Charge fee(s) indicated below, except for the filing fee to the above-identified deposit account.

FEE CALCULATION

1. BASIC FILING FEE

Large Entity	Small Entity	Fee Code	Fee Description	Fee Paid
1001 750	2001 375	Utility filing fee		
1002 330	2002 165	Design filing fee		
1003 520	2003 260	Plant filing fee		
1004 750	2004 375	Release filing fee		
1005 160	2005 80	Provisional filing fee		160.00
SUBTOTAL (1)				(\$)160.00

2. EXTRA CLAIM FEES FOR UTILITY AND REISSUE

Large Entity	Small Entity	Fee Code	Fee Description	Fee Paid
1202 18	2202 9	Claims in excess of 20		
1201 64	2201 32	Independent claims in excess of 3		
1203 280	2203 140	Multiple dependent claims, if not paid		
1204 64	2204 32	** Release independent claims over original patent		
1205 18	2205 9	** Release claims in excess of 20 and over original patent		
SUBTOTAL (2)				(\$)

**or number previously paid, if greater. For Release, see above

3. ADDITIONAL FEES

Large Entity	Small Entity	Fee Code	Fee Description	Fee Paid
1051 130	2051 65	Surcharge - late filing fee or oath cover sheet		
1052 50	2052 25	Surcharge - late provisional filing fee or cover sheet		
1053 130	2053 65	Non-English specification		
1812 2,520	2812 1,260	For filing a request for <i>ex parte</i> reexamination		
1804 920*	2804 460*	Requesting publication of SIF prior to Examiner action		
1805 1,840*	2805 920*	Requesting publication of SIF after Examiner action		
1251 110	2251 55	Extension for reply within first month		
1252 410	2252 205	Extension for reply within second month		
1253 930	2253 465	Extension for reply within third month		
1254 1,450	2254 725	Extension for reply within fourth month		
1255 1,870	2255 935	Extension for reply within fifth month		
1401 320	2401 160	Notice of Appeal		
1402 320	2402 160	Filing brief in support of an appeal		
1403 280	2403 140	Request for oral hearing		
1451 1,510	2451 755	Petition to institute a public use proceeding		
1452 110	2452 55	Petition to revive - unavoidable		
1453 1,300	2453 650	Petition to revive - unintentional		
1501 1,300	2501 650	Utility issue fee (for rollover)		
1502 470	2502 235	Design issue fee		
1503 630	2503 315	Plant issue fee		
1460 130	2460 65	Positions to the Commissioner		
1807 50	2807 25	Processing fee under 37 CFR 1.17(e)		
1808 180	2808 90	Submission of Information Disclosure Sheet		
8021 40	2021 20	Recording each patent assignment per property (does number of properties)		
1809 750	2809 375	Filing a submission after final rejection (37 CFR 1.129(b))		
1810 750	2810 375	For each additional invention to be examined (37 CFR 1.129(b))		
1801 750	2801 375	Request for Continued Examination (RCE)		
1802 800	2802 400	Request for expedited examination of a design application		

*Reduced by Basic Filing Fee Paid

SUBTOTAL (3) (\$)

SUBMITTED BY

Name (Print/Type) **Robin S. Quartin, Ph.D.**

Signature

Registration No. (Attorney/Agent) **45,028**

(Complete if applicable)

Telephone **302-885-9129**

Date **June 30, 2003**

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

This collection of information is required by 37 CFR 1.17 and 1.27. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (05-03)

Approved for use through 4/30/2003, OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number

100997-1/US

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle (if any))	Family or Surname	Residence (City and either State or Foreign Country)
Stewart	Fisher	Waltham, Massachusetts
Tom	Keating	Waltham, Massachusetts
Ann	Eakin	Waltham, Massachusetts
Steve	Marmor	Waltham, Massachusetts
Gunther	Kern	Waltham, Massachusetts
Elaina	Zverina	Waltham, Massachusetts

[Page 2 of 2]

Number 1 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

ISOTHERMAL TITRATION CALORIMETRY ASSAYS

5

FIELD OF THE INVENTION

10 The present invention relates to methods of identifying and characterizing modulators of bioprocesses and biomolecules.

BACKGROUND

15 A major process in drug discovery involves the study of biochemical systems and the specific nature of the interaction of biomolecules with various ligands. Microcalorimetric techniques such as isothermal titration calorimetry (ITC) and differential scanning calorimetry (DSC) are widely used in understanding these interactions (Jezesarov & Bosshard, 1999, J. Mol. Recognit., 12:3-18; Ward & Holdgate, 2001, Prog. Med. Chem., 38:309-76; Sturtevant, 1987, Ann. Rev. Phys. Chem., 38:463-88).

20 ITC is a technique used primarily in measuring the equilibrium heat of binding of a ligand to a macromolecule (Leavitt & Freire, 2001, Curr. Op. Struct. Biol., 11:560-6; Ladbury & Chowdhry, 1996, Chem. Biol., 3:791-801; Doyle, 1997, Biotechnology, 8:31-35; Fisher & Singh, 1995, Methods Enzymol., 259:194-221). In recent years ITC has also been used in the determination of enzymatic activity and enzyme kinetic parameters such as the
25 Michaelis-Menten kinetic parameters K_M and K_{cat} (Todd & Gomez, 2001, Anal. Biochem., 296:179-187; Cai *et al.*, 2001, Anal. Biochem., 299:19-23).

 Several studies of enzymatic activities have been reported using the detection of reaction heat for measuring kinetic parameters (Lonhienne *et al.*, 2001, Biochim. Biophys. Acta, 1545:349-356; Quemard *et al.*, 1995, Biochemistry, 34:8235-8241; Silberg & Vickery
30 2000, J. Biol. Chem., 275:7779-7786; Prodromou *et al.*, 1999, EMBO J., 18:754-762).

 The determination of the inhibition profiles of different compounds towards a target (*e.g.*, IC_{50} value, the concentration of compound at which 50% of the enzymatic activity is inhibited) is of particular interest in drug discovery. Determination of such quantities as IC_{50} and K_i (kinetically determined inhibition constant representing the potency of an enzyme
35 inhibitor) are imperative in designing novel potent inhibitors that will lead to developing

effective drugs. Recent experiments describe the use of DSC in determining K_d (equilibrium dissociation constant) values for binding of inhibitors to MurB (Sarver *et al.*, 2002, J. Biomol. Screening, 7:21-28). ITC is also used traditionally in the determination of K_d values but not IC_{50} or K_i values (kinetically determined inhibition parameters).

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic representation of a calorimetric chamber.

Figure 2A shows inhibition of *E. coli* GyrAB ATPase activity by novobiocin as measured by ITC. Figure 2B shows the data fitted to an IC_{50} curve.

10

Figure 3A shows *E. coli* GyrAB ATPase reaction progressing to completion after each addition of substrate. Figure 3B shows the heat calibration curve obtained by the same experiment.

Figure 4A shows inhibition of *E. coli* MurC activity by β,γ -Methyleneadenosine 5'-triphosphate as measured by ITC. Figure 4B shows the data fitted to IC_{50} curve.

15

Figure 5 shows inhibition of *E. coli* MurB activity by (2R)-2-[2-[3-(4-*tert*-butylphenoxy)phenyl]-4-oxo-1,3-thiazolidin-3-yl] hexanoic acid (data fitted to an IC_{50} curve).

Figure 6A shows the heat change $\Delta Q/\Delta T$ observed, in the presence (open circles) and absence of tetracycline (closed circles), upon titrating S30 extract (containing the ribosomes) in the reaction mixture (containing DNA, amino acids, and other necessary components for the reaction to occur). Figure 6B shows the difference in the heat change $\Delta(\Delta Q/\Delta T)$, observed in the presence and in the absence of tetracycline, upon titrating S30 extract in the reaction mixture.

25 DETAILED DESCRIPTION

The present invention provides methods for monitoring a bioprocess using isothermal titration calorimetry.

The present invention also provides methods for monitoring the activity of a biomolecule or a mixture of biomolecules using isothermal titration calorimetry. In some
30 embodiments, the biomolecule is an enzyme.

The present invention also provides methods for identifying and/or detecting modulators of a bioprocess by monitoring changes or alterations in the reaction heat produced by the bioprocess in the presence and in the absence of test modulators.

5 The present invention also provides methods for identifying and/or detecting modulators of a biomolecule by monitoring changes or alterations in the reaction heat produced by the activity or action of the biomolecule in the presence and in the absence of test modulators.

The present invention also provides methods for determining inhibition profiles (such as IC_{50} values) for compounds that inhibit a bioprocess

10 The present invention also provides methods for determining inhibition profiles (such as IC_{50} values) for compounds that inhibit a biomolecule. In some embodiments the biomolecule is an enzyme and an inhibition profile is determined for the activity of the enzyme.

15 The methods of the present invention have general applicability to kinetic measurements because the methods are based upon the determination of the heat of reaction, which is a universal quantity.

The methods of the present invention do not require coupling enzymes, and are devoid of artifacts and interferences that are sometimes produced in spectrophotometric and fluorometric assays.

20 The methods of the present invention are particularly useful for reactions where there is no straightforward way of detecting product formation or substrate depletion.

The present invention provides screening methods for identifying a biomolecule modulator. The methods comprise incubating a plurality of test mixtures, each test mixture comprising at least one biomolecule ligand or substrate, and at least one biomolecule, under
25 conditions effective to permit at least a portion of the biomolecule to react to a measurable extent; adding a test biomolecule modulator to at least one test mixture; detecting an activity signal of each biomolecule, as measured by a heat change, in the presence and in the absence of the biomolecule ligand or substrate; and comparing the activity signal of each biomolecule in the presence of the biomolecule modulator with an activity signal of the same biomolecule
30 in the absence of the biomolecule modulator under the same conditions.

In one embodiment of the invention the biomolecule is a protein.

In a particular embodiment of the invention the protein is an enzyme.

In a further particular embodiment of the invention the enzyme is gyrase (GyrAB), MurB, or MurC.

In a most particular embodiment of the invention the enzyme is *Escherichia coli* gyrase (GyrAB), *E. coli* MurB, or *E. coli* MurC.

In some embodiments of the present invention detecting the activity signal comprises detecting the change in energy input required to maintain the system's thermal equilibrium

- 5 In some embodiments, the detecting the activity signal comprises detecting the change in energy input required to maintain thermal equilibrium between the test mixture and a control.

- The present invention also provides methods for identifying a bioprocess modulator. The methods comprise incubating a plurality of test mixtures, each test mixture comprising at least one biomolecule having activity in the bioprocess, and at least one biomolecule ligand or substrate, wherein the conditions are effective to permit at least one biomolecule to react to a measurable extent; adding a test bioprocess modulator to at least one test mixture; detecting an activity signal of the bioprocess in the presence and in the absence of the biomolecule ligand or substrate; and comparing the activity signal of the bioprocess in the presence of the bioprocess modulator with a activity signal of the same biomolecule in the absence of the bioprocess modulator under the same conditions.
- 10
15

- Other embodiments of the invention comprise a method for identifying compounds that modulate a bioprocess comprising: measuring heat change in the bioprocess, in the presence and in the absence of a test compound; and identifying a compound that modulates the bioprocess as one that yields an alteration in heat change in the presence of the compound relative to heat change in the absence of the compound.
- 20

- Particular embodiments of the invention include methods for identifying compounds that modulate the activity of a biomolecule comprising: measuring heat change resulting from activity of the biomolecule, in the presence and in the absence of a test compound; and identifying a compound that modulates the biomolecule as one that yields an alteration in the heat change in the presence of the compound relative to the heat change in the absence of the compound.
- 25

- In addition, the methods of the present invention allow for the continuous measurement of initial rates ($\Delta Q/\Delta t$ can be measured instantly as the reaction occurs). Advantages of this are that problems such as product inhibition can be minimized in determining kinetic parameters of the reaction and/or inhibition profiles of potential inhibitors. Consequently, the methods of the present invention are particularly applicable to the screening and identification of novel modulators of diverse biomolecular processes.
- 30

Typically in ITC experiments, a macromolecule is placed in the sample cell, water or buffer is placed in the reference cell, and the cells are equilibrated; a preset differential power (DP) or baseline signal is applied to the cells such that ΔT is zero or constant (Figure 1). Upon reaching thermal equilibrium between the two cells, a ligand is injected into the sample cell through a titration syringe. Depending on the nature of the interaction between the macromolecule and the ligand, DP will decrease (exothermic event), or increase (endothermic) to compensate for the heat released or absorbed in the sample cell while keeping ΔT between the cells zero or constant.

In one aspect, the present invention provides methods for monitoring the inhibition of an enzyme, including methods of determining the inhibition profile of an inhibitor on an enzyme. In our approach, the cells are equilibrated after the enzymatic reaction is started and placed into the sample cell. The baseline signal DP is now lower than the preset value because the enzymatic reaction releases heat in the sample cell (exothermic reaction) in a steady state; consequently less power is needed to be applied to the sample cell to keep the two cells at the preset ΔT . During the experimental run, the inhibitor is injected into the sample cell. As the enzyme is inhibited, less heat is produced in the sample cell. The decrease in the heat produced by the enzymatic reaction per unit time results in an increase in the DP signal since more power must now be provided from the feedback heater to keep the two cells at the preset ΔT . The direction of the heat change would be opposite for an endothermic reaction.

The methods of the present invention are particularly useful for high throughput screening of enzyme inhibitors since the methods allow for the determination of the IC_{50} value in a single titration. There is no need to initiate the reaction after the system is equilibrated (the reaction is started before equilibration), and therefore there is no need for co-injection of a reaction initiator with the reaction inhibitor.

The methods of the present invention can be applied to the identification and characterization of enzyme modulators (inhibitors or activators) by a heat exchange assay. We have determined the kinetic and inhibition parameters for three different enzymatic systems. In general, monitoring of the change in heat with respect to time ($\Delta Q/\Delta t$) is used in measuring the initial rates of reactions under various conditions, including altered concentrations of enzyme, substrate, and modulator (inhibitor or activator). More specifically, we have characterized the enzymatic activity and studied modulation by known inhibitors of that enzymatic activity, for the following: the ATPase activity of *Escherichia*

coli gyrase (GyrAB), the redox activity of *E. coli* MurB (converts uridine diphosphate-*N*-acetyl-glucosamine enolpyruvate (EP-UDP-GlcNAc) to uridine diphosphate-*N*-acetylmuramic acid (UDP-MurNAc) with the simultaneous oxidation of NADPH to NADP⁺), and the ligase activity of *E. coli* MurC (converts UDP-MurNAc to uridine diphosphate-*N*-acetylmuramyl-L-alanine (UDP-MurNAc-Ala) with the simultaneous conversion of ATP to ADP and P_i).

As used herein, a "bioprocess" comprises one or more biomolecules, capable of catalyzing and/or supporting at least one reaction, in combination with, at a minimum, all of the necessary set of chemical reactants required for the reaction to occur. A bioprocess may comprise multiple reactions running simultaneously. Assay methods of the present invention can be used to monitor any or all of these reactions. Assay methods of the present invention can be used to screen for and identify compounds that modulate any or all of these reactions.

Assays of the present invention can be used to monitor any bioprocess that releases or absorbs heat. Assays of the present invention can be used to screen for compounds that modulate (inhibit or augment) any bioprocess.

As used herein, the terms "biomolecule" refer to chemical entities capable of catalyzing and/or supporting a chemical reaction in a bioprocess. Examples of biomolecules include, but are not limited to proteins, enzymes, polypeptides, DNA or RNA oligonucleotides, DNA or RNA polynucleotides, lipids, cofactors, and carbohydrates.

Assays of the present invention can be used to monitor the activity or action of any biomolecule-where, as a result of the activity or action, heat is released or absorbed. Assays of the present invention can be used to screen for compounds that modulate (inhibit or augment) the activity or action of any biomolecule.

In some embodiments, the bioprocess is an enzymatic reaction.

In some embodiments, the bioprocess is DNA supercoiling, the biomolecule is selected from type II topoisomerases including but not limited to DNA gyrase (GyrAB) and topoisomerase IV, and the chemical reactants are selected from adenosine triphosphate and DNA molecules including, but not limited to, oligonucleotides, PCR products, recombinant plasmids, cosmid, and chromosomal preparations.

In some embodiments, the ATPase activity of a DNA gyrase is measured as indicative of DNA supercoiling or unwinding.

In some embodiments, the ATPase activity of GyrAB measured as indicative of DNA supercoiling or unwinding.

In some embodiments, the bioprocess is a coupled process of chemical reactions comprising a metabolic pathway including, but not limited to, bacterial cell wall biosynthesis, aromatic amino acid biosynthesis, oxidative phosphorylation, citric acid cycle, glycolysis, gluconeogenesis, photophosphorylation, lipid biosynthesis, glycosylation and cofactor biosynthesis.

In some embodiments, the bioprocess is a coupled process of chemical reactions comprising a macromolecular process including, but not limited to, DNA replication, RNA polymerization, transcription/translation, protein synthesis, protein secretion, microtubule polymerization, histone deacetylation, proteasome protein degradation, chaperone mediated protein folding, and cellular respiration.

In some embodiments, the biomolecule is an enzyme.

Enzymes from any source can be monitored or modulated using the assays of the present invention, including, but not limited to, viral, bacterial, prokaryotic, eukaryotic, and cancer or disease-associated enzymes.

In some embodiments, the biomolecule is a bacterial enzyme.

In some embodiments, the biomolecule is a bacterial enzyme from *Escherichia coli*, *Salmonella spp*, *Shigella spp*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, *Chlamydia spp*, *Legionella spp*, *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Enterococcus faecium*, or *Mycoplasma spp*.

In some embodiments, the biomolecule is gyrase (GyrAB), MurB, or MurC.

In some embodiments, the biomolecule is *Escherichia coli* gyrase (GyrAB), *E. coli* MurB, or *E. coli* MurC.

In some embodiments of the present invention, the biomolecule is an enzyme selected from oxidases/reductases that utilize at least one of the following cofactors: NADH, flavin, cobalamin, S-adenosyl methionine, ubiquinone, heme, glutathione, photooxidation, and/or iron sulfur clusters.

In some embodiments, the biomolecule is the enzyme MurB and the chemical reactants are NADH, flavin, and UDP-GlcNAc-EP.

In some embodiments, the biomolecule is an ATP-dependent ligase, including, but not limited to, the enzymes FolC, MurC, MurD, MurE, MurF.

In some embodiments, the biomolecule is a kinase, including, but not limited to, protein serine/threonine kinases, protein tyrosine kinases, or dual specificity kinases.

In some embodiments, the biomolecule is a phosphatase, including, but not limited to, pyrophosphatase, protein phosphatases, and phosphodiesterases.

In some embodiments, the biomolecule is a transmembrane signaling protein, including, but not limited to, bacterial and eukaryotic two component regulatory systems, G-protein coupled receptor-G protein complexes, and JAK-STAT signaling complexes.

In some embodiments, the biomolecule is a protein chaperone, including, but not limited to, the heat shock proteins GroEL-GroES, DnaK.

The methods of the present invention utilize the measurement of heat change as an effect of a bioprocess. Heat change can be measured by a variety of techniques known to those of skill in the art.

In some embodiments of the present invention, heat change is measured using instruments designed for calorimetry. Such calorimeters, are known to the art and include commercially available instruments such as Microcal's isothermal titration calorimeter (Microcal VP ITC), and the differential scanning calorimeter (Microcal High-throughput DSC). In general, the activity of a bioprocess, in the presence and absence of test compounds and/or chemical reactants, is measured by monitoring the change in heat relative to a reference cell containing a control sample. Test compounds and/or chemical reactants can be introduced into sample and/or reference cells by various methods known to those skilled in the art, including, but not limited to, injection using one or more syringes.

In some embodiments, the sample and/or reference cell comprises a set of solution reservoirs on a thermal sensitive matrix, and the test compounds and/or chemical reactants are introduced using electrosmotic or vacuum-driven flow.

The sample cell can be in a variety of formats, including but not limited to, a section of capillary tubing, a set of solution reservoirs on a thermal sensitive matrix, a chamber enclosed in an adiabatic environment, etc.

The reference cell can be in a variety of formats, including but not limited to, a section of capillary tubing, a set of solution reservoirs on a thermal sensitive matrix, a chamber enclosed in an adiabatic environment, etc.

In some embodiments, heat change is measured using instruments designed for differential scanning calorimetry, and the heat change from chemical reactions resulting from a mixture of a test compound and a solution containing a bioprocess is measured in a sample cell over a temperature range relative to a control sample in a reference cell measured under the same conditions.

In some embodiments, the test compounds and/or chemical reactants are mixed and introduced into the sample cell using robotic liquid handling equipment.

In some embodiments, heat change is measured using a calorimeter selected from a Microcal ITC, a Microcal VP DSC, and a Microcal High throughput DSC.

5 The invention is further illustrated by way of the following examples, which are intended to elaborate several embodiments of the invention. These examples are not intended to, nor are they to be construed to, limit the scope of the invention. It will be clear that the invention may be practiced otherwise than as particularly described herein. Numerous modifications and variations of the present invention are possible in view of the teachings
10 herein and, therefore, are within the scope of the invention.

EXAMPLES

Example 1. Inhibition of gyrase by novobiocin as measured by ITC.

The enzyme comprising *E. coli* GyrA and GyrB subunits, at a concentration of 1 μ M
15 each, were incubated in 50 mM Tris (pH 7.5), 75 mM ammonium acetate, and 200 nM of 250
base pairs (bp) DNA at room temperature for 30 minutes. The enzyme stock was diluted 20-
fold to a final concentration of 50nM in assay buffer (50 mM Tris pH 7.5, 75 mM ammonium
acetate, 5% glycerol, 0.5 mM EDTA, 5.5 mM magnesium chloride) that contained ATP at a
final concentration of 500 μ M. The diluted enzyme solution was placed in the ITC sample
20 cell and the calorimetric cells were equilibrated. Novobiocin was diluted in assay buffer at a
final concentration of 2.8 μ M and it was loaded into the titration syringe. The titration was
started manually after the DP signal was stabilized. A total of 15 injections of 5 μ l each were
performed. Time spacing in between injections was set at 120 sec. Control experiments were
performed under identical conditions with experiment assay buffer in the syringe. In these
25 experiments, no decrease in DP baseline was observed after the completion of each injection
peak. In separate control experiments the linearity of the reaction was checked under the
same assay conditions. The reaction progress was found to be linear with time for at least 30
minutes. The inhibition curve of the ATPase activity of *E. coli* GyrAB by novobiocin, a
potent natural-product gyrase inhibitor, was generated by the ITC technique (see Figures 2A
30 and 2B). The reaction was started by adding ATP at a final concentration of 250 μ M to a
mixture of *E. coli* GyrAB and a 250 bp DNA. The final concentration of *E. coli* GyrAB in
the reaction was 50 nM of each subunit and the DNA concentration was 10nM. All
measurements contained 50 mM Tris, pH 7.5, 75 mM ammonium acetate, 5% glycerol, 0.5

mM EDTA, and 5.5 mM magnesium chloride. Novobiocin was dissolved in the same buffer at a final concentration of 2.8 μ M. The reaction was placed in the sample cell and the cells were equilibrated. Novobiocin was added in the titration syringe. Fifteen 5 μ l injections were performed.

5 Upon novobiocin addition, the power baseline decreased after each injection peak (solid black line in Figure 2A), an observation consistent with inhibition of enzyme activity. After a certain injection, the power baseline plateaus, an indication that all enzyme activity has been inhibited. The data were analyzed as follows: the power value (μ cal/sec) after each injection is subtracted from the initial power value (before any injection occurs). Each point
10 is then calculated as % inhibition assuming that 100% inhibition occurs where the power baseline plateaus after the n^{th} injection of inhibitor. After data analysis, an IC_{50} value of 26 nM was obtained for novobiocin inhibition. This value is in close agreement with the value obtained by using an independent assay of enzyme activity using phosphate detection. (IC_{50} = 28 nM, see Table 1).

15 It is also possible to convert the enzyme activity expressed as power units to activity expressed as turnover number units by using a calibration curve that converts $\Delta Q/\Delta t$ to $\Delta P/\Delta t$ and calculate the % inhibition as described above. An example of such conversion is shown in Figures 3A and 3B.

20 The experiment shown in Figure 3 was run as follows: reconstituted *E. coli* GyrAB (a mixture of 100 nM of each GyrA, and GyrB subunits in the presence of 20 nM of a 250bp DNA) was added into the sample cell. ATP was dissolved in the same buffer at a final concentration of 145 μ M and added to the titration syringe. Four injections (1, 3, 5, and 8 μ l) were completed. The injection spacings were adjusted so the power baseline returned to the initial position between injections, an indication that the reaction was completed for each
25 injection of substrate (see Figure 3A). The area under each peak represents the total heat released for the conversion of the added ATP to ADP and Pi. By plotting the concentration of ATP added in each injection vs. area of the injection peak it is possible to obtain a calibration curve the slope of which is the conversion factor $\Delta Q/\Delta P$ (heat released in μ cal/nmole of product formed). The slope of the curve shown in Figure 3B was calculated
30 to be 11kcal/mole. This number is consistent with the number of 13 kcal/mole reported for the energy released in ATP hydrolysis. Measuring kinetic parameters such as K_M , k_{cat} , and K_i , ADP by ITC contributed to further characterization of *E. coli* GyrAB. These values are

compared with those obtained by an independent assay of enzyme activity using phosphate detection (Table 1).

Table 1 shows kinetic parameters of the *E. coli* GyrAB ATPase reaction obtained by ITC. For comparison, values obtained by an independent method of phosphate detection are included.

Table 1

Method	$K_{M, ATP}$ (μM)	K_i, ADP (μM)	Novobiocin IC_{50} (μM)
ITC	166	18	0.026
Phosphate detection	145	18	0.028

Example 2. MurC.

E. coli MurC was diluted in assay buffer (50 mM Tris pH 8.0, 1 mM Tris (2-carboxyethyl) phosphine hydrochloride (TCEP), 0.01 % (w/v) Triton X-100, 20 mM ammonium formate) to a final concentration of 10 nM. The buffer also contained the MurC substrates at the following concentrations in the ITC sample cell: 100 μM UDP-MurNAc, 275 μM L-alanine, and 300 μM ATP. β , γ -Methyleneadenosine 5'-triphosphate (AMP-PCP) was dissolved in assay buffer at a final concentration of 2.8 mM and it was loaded into the titration syringe. The titration was started manually after the DP signal was stabilized. A total of 16 injections (2, 2, 4, 4, 4, 4, 8, 8, 16, 16, 32, 32, 32, 32, 32, and 32 μl) were performed. Time spacing in between injections was set at 120 sec for all injections.

The inhibition of MurC by AMP-PCP is shown in Figure 4. All the reactions were carried in 50 mM Tris, pH 8.0, 1 mM TCEP, 0.01 % (w/v) Triton, and 20 mM ammonium formate. The final concentration of *E. coli* MurC was 10 nM. The final concentrations of the MurC substrates were as follows: 100 μM uridine diphosphate-N-acetylmuramic acid (UDP-MurNAc), 275 μM L-alanine, and 300 μM ATP. The reaction was initiated with the addition of MurC and placed into the sample cell. AMP-PCP was dissolved in the buffer described above at a final concentration of 2.8 mM and placed into the titration syringe.

The raw ITC data are shown in Figure 4A. An IC_{50} value of 27 μM was obtained for MurC inhibition by AMP-PCP (see Figure 4B). This value compared well with the value of 40 μM obtained using an independent assay of enzyme activity by detecting inorganic phosphate formation.

Other kinetic measurements were performed to further characterize *E. coli* MurC by ITC. Table 2 shows kinetic parameters of the *E. coli* MurC reaction obtained by ITC. For comparison, values obtained by an independent method of phosphate detection are included.

Table 2

Method	K_M, ATP (μM)	$K_M, UNAM$ (μM)	K_M, ALA (μM)	k_{cat} (min^{-1})	AMP-PCP IC_{50} (μM)
ITC	61	21	44	50	27
Phosphate detection	101	19	56	71	35

5

Example 3. MurB.

E. coli MurB was diluted in assay buffer (50 mM Tris pH 8.0, 1 mM TCEP, 20 mM KCl, 0.005 % Triton X-100) to a final concentration of 1.3 nM. The buffer contained the MurB substrates at the following concentrations in the ITC sample cell: 100 μM NADPH and 60 μM EP. DMSO was also added at a final concentration of 2.7 % (v/v). The compound ((2R)-2-[2-[3-(4-*tert*-butylphenoxy)phenyl]-4-oxo-1,3-thiazolidin-3-yl] hexanoic acid) stock was prepared in DMSO and diluted in assay buffer at a final concentration of 2.8 mM. The titration was started manually after the DP signal stabilized and a total of 10 injections (8, 8, 17, 17, 24, 24, 32, 32, and 50 μl) were performed. Time spacing in between injections was set at 100, 100, 120, 120, 120, 150, 150, 150, 150, and 180 sec respectively.

15

The inhibition of *E. coli* MurB by (2R)-2-[2-[3-(4-*tert*-butylphenoxy)phenyl]-4-oxo-1,3-thiazolidin-3-yl] hexanoic acid (a 4-thiazolidinone compound) was studied by the same method. Reactions were carried in 50 mM Tris, pH 8.0, 1 mM TCEP, 20 mM KCl, 0.005 % Triton, in the presence of 100 μM NADPH and 60 μM EP. The concentration of MurB was 1.3 nM. An IC_{50} value of 154 μM was obtained by the method described above (see Figure 5). An IC_{50} value of 101 μM was obtained by an absorbance method of measuring MurB activity by measuring NADPH depletion at 340 nm.

20

Other kinetic measurements were performed to further characterize *E. coli* MurB by ITC. Table 3 shows kinetic parameters of the *E. coli* MurB reaction obtained by ITC. For comparison, values obtained by an independent method of NADPH depletion detection are included.

25

Table 3

Method	$K_M, NADPH$ (μM)	K_M, EP (μM)	4-thiazolidinone analogue IC_{50} (μM)
--------	-----------------------------	--------------------------	---

ITC	9	8	154
Absorbance	9	6	101

Example 4. Inhibition of the transcription/translation process by tetracycline as measured by ITC.

The inhibition of the transcription/translation process by tetracycline is shown in figure 6.

- 5 A reaction mix was prepared, containing 17.5 mM Tris acetate, 95.2 mM potassium acetate, 15 mM ammonium acetate, 5mM TCEP, 5 mM ATP, 1.25 mM each of four nucleotide triphosphates, adenosine triphosphate (ATP), cytidine triphosphate (CTP), guanine triphosphate (GTP), and uridine triphosphate (UTP), 87 mM phospho(enol)pyruvate (PEP), 0.25 mg/ml transfer ribonucleic acid (tRNA), 350 mg polyethyleneglycol (PEG), 0.05 mg/ml
- 10 folinic acid, 2.5 mM cyclic adenosine monophosphate (cAMP), 2mM isopropyl b-D-1-thiogalactopyranoside (IPTG), and 18.9 mM each of the 20 amino acids.

A 2 ml substrate mixture was prepared, containing 800 μ l of the reaction mixture, 15 μ l of 1 mg/ml pBEST *luc*TM vector DNA (Promega) and 1185 μ l H₂O.

- The same substrate mixture was prepared, containing containing 800 μ l of the reaction mixture, 15 μ l of 1 mg/ml pBEST *luc*TM vector DNA (Promega), 20 μ l of 20 mM tetracycline in dimethyl sulfoxide (DMSO), and 1165 μ l H₂O.
- 15

- S30 extract (containing ribosomes and a wide variety of enzymes and cofactors needed for the transcription/translation process) from bacterial lysate was prepared and added in the titration syringe. S30 extract was titrated in the sample cell containing the substrate mixture. 10 injections, 10 μ l each were performed Initially upon S30 extract addition the power baseline increased after each injection (indicative of an endothermic event). At the fifth injection of S30 extract the power baseline decreased and continued decreasing after subsequent injections (indicative of exothermic event) (see Figure 6A, closed circles).
- 20

- S30 extract was titrated in the sample cell containing the substrate mixture and tetracycline (see Figure 6A, open circles). Initially the same heat changes were observed, but at the fifth injection of S30 extract (exothermic event starts to take place), differences in the magnitude of the heat change were evident. These differences were calculated and plotted as differences in the heat change observed $\Delta(\Delta Q/\Delta T)$ upon titrating S30 extract in the substrate mixture in the presence or absence of tetracycline (see Figure 6B). $\Delta(\Delta Q/\Delta T)$ is defined as the heat change $\Delta Q/\Delta T$ observed in the presence of tetracycline minus the heat change
- 25
- 30

- 5 $\Delta Q/\Delta T$ observed in the absence of tetracycline. It is evident from figure 6B that a portion of the transcription/translation process is inhibited by tetracycline since at the fifth injection of S30 extract the heat change is smaller in the presence of tetracycline (negative $\Delta(\Delta Q/\Delta T)$) than that in the absence of tetracycline. The effect of tetracycline in the heat change of the process becomes less evident upon continuing S30 injections (the differences in the heat change observed in the presence and absence of tetracycline become smaller, or $\Delta(\Delta Q/\Delta T)$ returns to 0).

- 10 The foregoing examples are meant to illustrate the invention and are not to be construed to limit the invention in any way. Those skilled in the art will recognize modifications that are within the spirit and scope of the invention.

We claim:

1. A screening method for identifying a biomolecule modulator, the method comprising:
 - a) incubating a plurality of test mixtures, each test mixture comprising at least one biomolecule ligand or substrate, and at least one biomolecule, under conditions effective to permit at least a portion of the biomolecule to react to a measurable extent;
 - 5 b) adding a test biomolecule modulator to at least one test mixture;
 - c) detecting an activity signal of each biomolecule as measured by a heat change in the presence and in the absence of the biomolecule ligand or substrate; and
 - d) comparing the activity signal of each biomolecule in the presence of the biomolecule modulator with a activity signal of the same biomolecule in the absence of the biomolecule modulator under the same conditions.
- 10 2. The method of claim 1 wherein the biomolecule is a protein.
3. The method of claim 1 wherein the biomolecule is an enzyme.
- 15 4. The method of claim 3 wherein the enzyme is gyrase (GyrAB), MurB, or MurC.
5. The method of claim 3 wherein the enzyme is *Escherichia coli* gyrase (GyrAB), *E. coli* MurB, or *E. coli* MurC.
- 20 6. The method of claim 4 wherein the enzyme is GyrAB.
7. The method of claim 6 further wherein ATPase activity is measured.
- 25 8. The method of claim 1 wherein detecting the activity signal comprises detecting the change in energy input required to maintain thermal equilibrium between the test mixture and a control.
9. The method of claim 1, adapted for high throughput screening.
- 30 10. The method of claim 1, wherein the method is carried out in a calorimeter.
11. A screening method for identifying a bioprocess modulator, the method comprising:

- a) incubating a plurality of test mixtures, each test mixture comprising at least one biomolecule having activity in the bioprocess, and at least one biomolecule ligand or substrate, wherein the conditions are effective to permit at least one biomolecule to react to a measurable extent;
- 5 b) adding a test bioprocess modulator to at least one test mixture;
- c) detecting an activity signal of the bioprocess in the presence and in the absence of the biomolecule ligand or substrate; and
- d) comparing the activity signal of the bioprocess in the presence of the bioprocess modulator with a activity signal of the same biomolecule in the absence of the bioprocess modulator under the same conditions.
- 10
12. The method of claim 11 wherein the bioprocess is transcription/translation.
13. A method for identifying compounds that modulate a bioprocess comprising:
- 15 a) measuring heat change in the bioprocess, in the presence and in the absence of a test compound; and
- b) identifying a compound that modulates the bioprocess as one that yields an alteration in heat change in the presence of the compound relative to heat change in the absence of the compound.
- 20
14. A method for identifying compounds that modulate the activity of a biomolecule comprising:
- a) measuring heat change resulting from activity of the biomolecule, in the presence and in the absence of a test compound; and
- 25 b) identifying a compound that modulates the biomolecule as one that yields an alteration in heat change in the presence of the compound relative to heat change in the absence of the compound.

ABSTRACT

An isothermal analysis method for measuring the activity of modulators of biomolecules.

FIGURE 1




FIGURE 2A




FIGURE 2B




FIGURE 3A




FIGURE 3B




FIGURE 4A




FIGURE 4B




FIGURE 5




FIGURE 6A




FIGURE 6B

